

Unruptured Brain Aneurysm

Overview

An aneurysm is a balloon-like bulge of an artery wall. As an aneurysm grows it puts pressure on nearby structures and may eventually rupture. Most people find out they have an unruptured aneurysm by chance during a scan for some other problem. The risk of rupture varies depending on the aneurysm location and size. Treatment options include observation, clipping, coiling, flow diversion, and bypass.

Blood supply of the brain

To understand aneurysms, it is helpful to know how blood circulates to the brain (see Anatomy of the Brain). Blood is pumped from the heart and carried to the brain by two paired arteries, the internal carotid arteries and the vertebral arteries (Fig. 1). The internal carotid arteries supply the anterior (front) areas and the vertebral arteries supply the posterior (back) areas of the brain. After passing through the skull, the right and left vertebral arteries join together to form a single basilar artery. The basilar artery and the internal carotid arteries communicate with each other in a ring at the base of the brain called the Circle of Willis.

What is an unruptured aneurysm?

An aneurysm is a balloon-like bulge or weakening of an artery wall. (Similar to a balloon on the side of a garden hose.) As the bulge grows it becomes thinner and weaker. It can become so thin that the blood pressure within can cause it to leak or burst open — a life-threatening hemorrhage in the brain. The vast majority of aneurysms are silent, meaning they have no symptoms until they rupture. It is impossible to predict if and when an aneurysm may rupture, but when they do it can cause major disability and be fatal in 40% of people. On the other hand, many people die of old age with an aneurysm, but not because of it. So, not every aneurysm is at high risk of rupture.

Aneurysms usually occur on larger blood vessels at the fork where an artery branches off. Types of aneurysms include (Fig. 2):

- **Saccular** - (most common, also called "berry") the aneurysm bulges from one side of the artery and has a distinct neck at its base.
- **Fusiform** - the aneurysm bulges in all directions and has no distinct neck.




Figure 1. The internal carotid arteries form the anterior (green) circulation and the vertebral / basilar arteries form the posterior (red) circulation. About 80% of aneurysms arise in the anterior system; 20% form in the posterior. Most aneurysms arise on the Circle of Willis.




Figure 2. A saccular aneurysm arises at the weak point of the artery wall where it branches. Other aneurysm types include dissecting and fusiform.

- **Dissecting** – a tear in the inner wall of the artery allows blood to split the layers and pool; often caused by a traumatic injury.

Aneurysms are also classified by size:

- Small = less than 7 millimeters in diameter
- Medium = 7-12 millimeters
- Large = 13-24 millimeters (size of a dime)
- Giant = more than 25 millimeters (quarter size)

The risk of aneurysm rupture is about 1% but may be higher or lower depending on the size and location of the aneurysm. Generally, the larger the aneurysm (>12mm), the higher risk of rupture. Also, aneurysms in the posterior circulation (basilar, vertebral and posterior communicating arteries) have a higher risk of rupture.

Risk factors for rupture include smoking, high blood pressure, drug or alcohol abuse, atherosclerosis, and genetic factors (1, 2). Lifestyle changes can reduce your risk of rupture.

What are the symptoms?

Most aneurysms don't have symptoms until they rupture. Ruptured aneurysms release blood into the spaces around the brain called a subarachnoid hemorrhage (SAH). Unruptured aneurysms rarely show symptoms until they grow large or press on the brain or nerves. Rupture usually occurs while a person is active rather than asleep. If you experience the symptoms of a SAH, call 911 immediately!

Symptoms of a ruptured aneurysm / SAH

- Sudden onset of a severe headache (often described as "the worst headache of my life")
- Nausea and vomiting
- Stiff neck
- Transient loss of vision or consciousness

Symptoms of an unruptured aneurysm

- Double vision
- Dilated pupils
- Pain above and behind the eye
- Newly unexplained headaches (rare)

What are the causes?

Studies have shown a strong link to family history (2). If an immediate family member has suffered an aneurysm, you are 4 times more likely to have one as well. The genetic link is not completely understood and studies are underway to determine if there is a pattern of inheritance. The most important inherited conditions associated with aneurysms include Ehlers-Danlos IV, Marfan's syndrome, neurofibromatosis NF1, and polycystic kidney disease. For those with a family history of aneurysm in two or more blood relatives, we recommend a screening test with a CT or MR angiogram.

Who is affected?

About 2 to 3% of Americans may have or develop an aneurysm; of those, 15% have multiple aneurysms. Unruptured aneurysms are more common than ruptured (1). However, 85% of aneurysms are not diagnosed until after they bleed. Aneurysms are usually diagnosed between ages 35 to 60 and are more common in women.

How is a diagnosis made?

Most people find out they have an unruptured aneurysm by chance (incidental) during a scan for some other medical problem. If you are experiencing symptoms and your primary care doctor suspects an aneurysm, you may be referred to a neurosurgeon. The surgeon will learn as much about your symptoms, current and previous medical problems, current medications, family history, and perform a physical exam. Diagnostic tests are used to determine the aneurysm's location, size, type, and involvement with other structures.

- **Computed Tomography (CT)** is a noninvasive X-ray to review the anatomical structures within the brain and to detect bleeding in or around the brain. CT angiography involves the injection of contrast into the blood stream to view the arteries of the brain.
- **Angiogram** is an invasive procedure, where a catheter is inserted into an artery and passed through the blood vessels to the brain. Once the catheter is in place, a contrast dye is injected into the bloodstream and the x-ray images are taken.
- **Magnetic Resonance Angiography (MRA)** is a noninvasive test, which uses a magnetic field and radio-frequency waves to give a detailed view of the soft tissues of your brain. An MRA involves the injection of contrast into the blood stream to examine the blood vessels, as well as the soft tissues of the brain.

What treatments are available?

Deciding how, or even if, to treat an unruptured aneurysm involves weighing the risks of rupture versus the risks of treatment. In addition to the size and location of the aneurysm, your overall health and medical history must be considered. The neurosurgeon will discuss with you all the options and recommend a treatment that is best for your individual case.

Observation

Sometimes the best treatment may be to simply watch the aneurysm over time and reduce your risk of rupture. You should quit smoking and control high blood pressure. Aneurysms that are small, unruptured, and asymptomatic may be observed with imaging scans every year until the growth or symptoms necessitate surgery. Observation may be the best option for patients with other health conditions.

Surgical clipping

Clipping is an open surgery performed under general anesthesia. An opening is cut in the skull, called a craniotomy. The brain is gently retracted so that the artery with the aneurysm may be located. A small clip is placed across the neck of the aneurysm to block the normal blood flow from entering the aneurysm (Fig. 3). The clip is made of titanium and remains on the artery permanently. Recovery time typically is four to six weeks, but symptoms may last longer.

Artery occlusion and bypass

If the aneurysm is large and inaccessible or the artery is too damaged, the surgeon may perform a bypass surgery. A craniotomy is cut to open the skull and clips are placed to completely block off (occlude) the artery and aneurysm. The blood flow is then rerouted (bypassed) around the occluded artery by inserting a vessel graft. The graft is a small artery, usually taken from your leg, which is connected above and below the blocked artery so that blood flows through the graft.

Another method is to detach a donor artery from its normal position on the scalp and connect it above the blocked artery inside the skull. This is called a STA-MCA (superficial temporal artery to middle cerebral artery) bypass. Recovery time typically is four to six weeks, but symptoms may last longer.

Endovascular coiling

In contrast to open surgery, a minimally invasive option is endovascular coiling. It is performed during an angiogram in the radiology department. During a coiling procedure, a catheter is inserted into an artery in the groin and then passed through the blood vessels to the aneurysm in the brain. The doctor guides the catheter through the bloodstream while watching an x-ray monitor. Through the catheter, the aneurysm is packed with a basket of platinum coils. The coils induce clotting (embolization), which seals off the aneurysm and prevents blood from entering (Fig. 4). Recovery time typically is two to four days. Follow-up imaging is performed periodically for 5 years to confirm the coils have not compacted and the aneurysm is not regrowing.

Endovascular flow diversion

If clipping or coiling would be difficult due to the shape, size, or wide neck of the aneurysm, a flow diversion stent may be used. A flow-diverter is a tightly woven mesh tube placed inside the parent artery across the aneurysm (Fig. 5). Because blood cannot easily get through the spaces of the tight mesh stent, the blood flows inside the flow-diverter and continues down the artery without going into the aneurysm. Without the pulsating blood flow, the aneurysm will eventually clot off and shrink. Recovery time typically is two to four days.




Figure 3. A titanium clip is placed across the neck of an aneurysm so that the blood flows through the artery, not the aneurysm.




Figure 4. The aneurysm is packed with platinum coils. The coils induce clotting (embolization), which seals off the space and prevents blood from entering into the aneurysm.




Figure 5. A flow-diverter mesh stent is placed inside the artery to reduce blood flow from entering the aneurysm. The aneurysm will eventually clot off and shrink.

Clinical trials

Clinical trials are research studies in which new treatments—drugs, diagnostics, procedures, and other therapies—are tested in people to see if they are safe and effective. Research is always being conducted to improve the standard of medical care. Information about current clinical trials, including eligibility, protocol, and locations, are found on the Web. Studies can be sponsored by the National Institutes of Health (see www.clinicaltrials.gov) as well as private industry and pharmaceutical companies (see www.centerwatch.com).

Should the aneurysm be treated?

Deciding how, or even if, to treat an unruptured aneurysm involves weighing the risks of rupture versus the risks of treatment. In general, surgical craniotomy carries a higher risk of brain disability in cognition and memory, especially for people over age 65. But endovascular procedures have a higher risk of aneurysm recurrence. Listed are general pros and cons to discuss with your neurosurgeon:

	Pros	Cons
Observation	No risk from intervention Non-invasive MRA scans confirm aneurysm size is stable	Risk of rupture variable based on size, location, age, smoking, hypertension, and family history
Surgical craniotomy (clip, bypass)	95% lifetime guarantee won't regrow 99% lifetime guarantee won't rupture	Risk of brain disability (memory, thinking) from craniotomy Risk of seizure
Endovascular procedure (coil, flow diversion)	Lower risks of brain disability (memory, thinking) than surgery 80-85% lifetime guarantee won't regrow 98% lifetime guarantee won't rupture	Risk of regrowth due to coil compaction Invasive follow-up angiograms at 6, 12 and 24 months

Recovery & prevention

Unruptured aneurysm patients recover from surgery or endovascular treatment much faster than those who suffer a SAH. Patients may suffer short-term and/or long-term deficits as a result of aneurysm treatment or rupture. Some of these deficits may disappear over time with healing and therapy.

Sources & links

If you have more questions, please contact Mayfield Brain & Spine at 800-325-7787 or 513-221-1100.

Links

The Brain Aneurysm Foundation
www.bafound.org 617-723-3870

www.brainaneurysm.com

Sources

1. Wiebers DO: Unruptured intracranial aneurysms – risk of rupture and risks of surgical intervention. *N Engl J Med* 339:1725-33, 1998.
2. Thompson BG: Guidelines for the management of patients with unruptured intracranial aneurysms. *Stroke* 46:2368-2400, 2015
3. Rates of delayed rebleeding from intracranial aneurysms are low after surgical and endovascular treatment. *Stroke* 37:1437-42, 2006

Glossary

aneurysm: a bulge or weakening of an artery wall.

coiling: a procedure to insert platinum coils into an aneurysm; performed during an angiogram.

Ehlers-Danlos IV: a genetic disorder of the connective tissue in the intestines, arteries, uterus, and other hollow organs that may lead to organ or blood vessel rupture.

embolization: inserting material, coil or glue, into an aneurysm so blood can no longer flow through it.

Marfan's syndrome: a genetic disorder in which patients develop skeletal defects in long bones, chest abnormalities, curvature of the spine, and circulatory defects.

neurofibromatosis (NF1): a genetic disorder, also called von Recklinghausen disease, in which patients develop café-au-lait spots, freckling, and multiple soft tumors under the skin and throughout the nervous system.

polycystic kidney disease: a genetic disorder in which patients develop multiple cysts on the kidneys; associated with aneurysms of blood vessels in the brain.

subarachnoid hemorrhage: bleeding in the space surrounding the brain; may cause a stroke.



updated > 1.2020

reviewed by > Andrew Ringer, MD and Craig Kilburg, MD, Mayfield Clinic, Cincinnati, Ohio

Mayfield Certified Health Info materials are written and developed by the Mayfield Clinic. We comply with the HONcode standard for trustworthy health information. This information is not intended to replace the medical advice of your health care provider. © Mayfield Clinic 1998-2020.