

## Percutaneous Stereotactic Radiofrequency Rhizotomy

### Overview

Percutaneous stereotactic radiofrequency rhizotomy (PSR) is a minimally invasive procedure performed to relieve pain caused by trigeminal neuralgia, glossopharyngeal neuralgia, and cluster headaches. Medications often provide initial pain relief, but may become ineffective or cause serious side effects. PSR uses a heating current to destroy some of the nerve fibers carrying pain signals to the brain. The patient will now feel some numbness in the area rather than pain.

### What is PSR?

Percutaneous stereotactic rhizotomy (PSR) can relieve neuralgia (nerve pain) by destroying the part of the nerve that causes pain and by suppressing the pain signal to your brain. The surgeon passes an electrode introducer (hollow needle) through the skin of your cheek into the selected nerve at the base of the skull. A heating current, which is passed through the electrode, destroys some of the nerve fibers (Fig. 1). The entire nerve is not destroyed. However, damaging the nerve causes mild to major facial numbness in that area. A degree of facial numbness is an expected outcome of the procedure and is necessary to achieve long-term pain relief. PSR is used to treat:

- **trigeminal neuralgia:** an irritation of the fifth cranial nerve causing severe pain that usually affects one side of the face, normally in the forehead, cheek, jaw, or teeth.
- **glossopharyngeal neuralgia:** an irritation of the ninth cranial nerve causing pain on one side of the throat near the tonsil area that can radiate to your ear and the angle of the jaw.
- **cluster headache:** attacks of severe pain on one side of the head behind the eye that last 30 to 90 minutes and recur one or more times a day at the same time usually for 4 to 12 weeks.

### Who is a candidate?

PSR can provide pain relief for many patients with trigeminal neuralgia, glossopharyngeal neuralgia, or other neurological diseases (e.g., cluster headache) when medications become ineffective. PSR can be effective in treating patients of all ages including those with multiple sclerosis and those with some types of tumors. Patients often choose to undergo PSR because it poses lower surgical risks than those




Figure 1. Trigeminal neuralgia is extreme, shock-like pain caused by faulty firing of the nerve. During PSR, an electrode is passed along the cheek through a hole in the skull to reach the trigeminal ganglion. Once the exact area triggering pain is located, a heating current selectively destroys some of the nerve fibers.

of a major operation such as microvascular decompression (MVD). Although facial numbness results, PSR is one of the most effective procedures because it provides lasting pain relief with few risks of serious side effects.

### Who performs the procedure?

PSR is performed by a neurosurgeon as an outpatient procedure in a radiology department or an operating room.

### What happens before treatment?

The surgeon will explain the procedure, its risks and benefits, and you will have time to ask questions. Consent forms are signed and paperwork completed to inform the surgeon about your medical history (e.g., allergies, medicines, anesthesia reactions, previous surgeries). Presurgical tests (e.g., blood test, electrocardiogram) may need to be done several days before surgery. Consult your primary




Figure 2. A nurse delivers anesthesia to a patient during PSR. The surgeon will adjust the current that destroys the pain-causing nerve fibers on a radiofrequency generator (A). The x-ray monitor (B) shows a view of the brain from the fluoroscope (C). The patient's arms are secured onto the narrow table and a grounding wire (D) is attached.

care physician about stopping certain medications and ensure you are cleared for surgery.

Stop taking all non-steroidal anti-inflammatory medicines (ibuprofen, naproxen, etc.) and blood thinners (Coumadin, aspirin, Plavix, etc.) 7 days before surgery.

You may be asked to wash your skin and hair with Hibiclens (CHG) or Dial soap before surgery. It kills bacteria and reduces surgical site infections. (Avoid getting CHG in eyes, ears, nose or genital areas.)

Patients are admitted to the outpatient clinic several hours before the procedure. No food or drink is permitted past midnight the night before undergoing PSR. The surgeon will give you specific instructions about taking your medications the day of the procedure.

### What happens during treatment?

There are 5 steps of the procedure, which generally takes 1 to 2 hours.

#### Step 1: prepare the patient

Several points are marked with a pen on your cheek, which is then washed with an antiseptic. Sterile drapes cover your neck and chest. You will receive a shot of atropine to help dry your mouth. A nurse assists the surgeon during PSR in checking your blood pressure and heart rate and delivering medications through an IV in your arm (Fig. 2).

#### Step 2: insert the electrode

As short-acting anesthesia is given, you will feel a slight burning sensation as you fall asleep. For trigeminal neuralgia, the surgeon guides the electrode introducer through your cheek into an opening at the base of the skull, called the foramen ovale, where the mandibular division (V3) of the trigeminal nerve exits the skull (Fig. 3A). Correct placement of the introducer is verified by x-ray.

For glossopharyngeal neuralgia, the surgeon inserts the introducer into the skull base at the jugular foramen where the glossopharyngeal nerve exits.





Figure 3. (top) The surgeon guides the electrode introducer through the cheek, into the foramen ovale, and beyond the ganglion to reach the nerve root fibers. (bottom) By advancing the electrode and rotating the curved tip, the surgeon locates the specific division of the trigeminal nerve causing pain.

### Step 3: find the source of your pain

During this step, you are awakened to help the surgeon identify in which trigeminal division the pain occurs. A low current of electricity, stimulating the nerve, reproduces the location of the pain. Based on your feedback, the surgeon inserts the electrode so that tingling occurs where your painful attacks are located (Fig. 3B).

### Step 4: destroy the pain-causing fibers

You will receive a mild anesthetic again to block any pain or discomfort. While you are asleep, the surgeon passes a heating current through the electrode. This current destroys the portion of the nerve that transmits pain and disrupts the pain-producing signal (Fig. 4).

### Step 5: test for pain or numbness

You are awakened to help the surgeon determine if the heating current has disrupted the pain. Areas of your face are tested for sensitivity by needle prick. Based on the amount of pain or numbness you feel, the surgeon determines how much of the nerve should be treated. Steps 3 through 5 may be repeated until the desired pain relief is achieved.

### What happens after treatment?

After the procedure, you will recuperate in a hospital room for 4 to 6 hours. An ice pack, placed on your jaw, reduces swelling caused by the operation. Patients go home the same day.

While adjusting to the numbness, you should be careful when shaving, eating/drinking hot foods/liquids, and chewing. A soft diet is often recommended for the first few weeks. Dentures, if used, can be worn at any time. Patients taking anticonvulsant or pain medication for trigeminal neuralgia prior to PSR will be weaned off the medications according to a schedule to decrease risk of withdrawal and side effects.

### Discharge instructions:

- Do not rub your eyes. Eye drops may be prescribed. Please use as directed. Inspect your eye daily for signs of redness, irritation, or blurring of vision. Notify your surgeon if this occurs.
- It will take time to adjust to the numbness in your face. Follow these tips:
  - Avoid very hot or extremely cold foods
  - Be careful while shaving
  - Don't bite the side of your lip where it is numb
- Mouth care is important. Use dental floss, rinse after meals, and watch for any signs of infection or irritation inside your cheek region.
- Swallowing may be an issue for a few days due to jaw pain. Eat a soft foods diet for 2 weeks and chew slowly.




Figure 4. Short controlled bursts of heat are applied to selectively destroy portions of the nerve.

### What are the results?

For trigeminal neuralgia, 98% of patients have immediate pain relief after PSR. About 20% of those who undergo PSR experience some recurrence of pain within 15 years; about half of these patients will require medication or undergo another procedure to control pain.

For glossopharyngeal neuralgia, 90% of patients have immediate pain relief after PSR. In patients with vagoglossopharyngeal neuralgia from neck tumors, 70% of patients can expect pain relief or improvement after PSR. Open rhizotomy is the treatment of choice for patients who experience vagoglossopharyngeal syncope.

For cluster headache, almost all patients have some pain relief after PSR. Only 62% of those experience long-term pain relief and will require medication or another procedure to control pain. PSR works best in treating pain behind the eye, but is not as effective in treating pain that radiates to the temple and ear region. Regardless of the surgery that is performed, good pain control rather than complete pain relief is expected.

### Recurrence

If pain recurs, inform your doctor. Excellent results can be achieved by percutaneous procedures, especially PSR, with less risk of operative complications. Healthy patients with pain recurring after a prior percutaneous procedure are best treated by microvascular decompression (MVD). A repeat percutaneous procedure or radiosurgery is



considered for patients who are medically unhealthy. Patients who have recurrent trigeminal neuralgia in an already numb area rarely benefit from repeat percutaneous surgery, but pain may be relieved by microvascular decompression or complete section of the sensory and motor root (neurectomy).

### What are the risks?

For trigeminal neuralgia, numbness, the most common side effect of PSR, is necessary for pain relief. Numbness usually occurs in the cheeks, gums, teeth, or tongue. Dysesthesia (troublesome numbness) is reported in 7% of patients after PSR. The loss of corneal reflex is reported in 6% of PSR procedures, primarily in patients with V-1 nerve pain. Complications (e.g., bleeding or infection) are rare. Other possible side effects include blurred or double vision, as well as weakness in the jaw (e.g., making chewing difficult); these symptoms usually resolve within 6 months. Some patients may develop fever blisters that will heal in 1–2 weeks.

For glossopharyngeal neuralgia, common side effects are hoarseness, vocal cord paralysis, and dysphagia (difficulty in swallowing). Controlled PSR for ninth and tenth cranial nerve neuralgia is more difficult than for trigeminal neuralgia. PSR should be restricted to patients with cancer pain who already have swallowing problems and vocal cord paralysis.

For cluster headache, because the nerve pain originates in V1, major sensory loss is associated with increased risks of keratitis and dysesthesia that can be up to 12%. For this reason, balloon compression may be worthy of consideration for the treatment of cluster headache.

### Sources & links

If you have more questions, please contact Mayfield Brain & Spine at 800-325-7787 or 513-221-1100.

### Support

Support groups provide an opportunity for patients and their families to share experiences, receive support, and learn about advances in treatments, pain control, and medications. Please contact the TNA Facial Pain Association at 800-923-3608.

### Sources

- 1) Taha JM, Tew JM Jr, Buncher CR: A prospective 15-year follow-up of 154 consecutive patients with trigeminal neuralgia treated by percutaneous stereotactic radiofrequency rhizotomy. *J Neurosurg* 83:989-993, 1995.

- 2) Taha JM, Tew JM Jr: Comparison of surgical treatments for trigeminal neuralgia: reevaluation of radiofrequency rhizotomy. *Neurosurgery* 38:865-871, 1996.
- 3) Taha JM, Tew JM Jr: Long-term results of radiofrequency rhizotomy in the treatment of cluster headache. *Headache* 35:193-196, 1995.

### Links

TNA Facial Pain Association [www.fpa-support.org](http://www.fpa-support.org)

### Glossary

**diplopia:** double vision.

**dysesthesia:** troublesome or bothersome numbness.

**dysphagia:** difficulty in swallowing.

**electrode:** a conductor that carries current. It is passed through a hollow needle called the electrode introducer.

**fluoroscope:** uses x-ray or other radiation to view structures in the body.

**ganglion:** a group of nerve cell bodies.

**keratitis:** inflammation of the cornea.

**neuralgia:** severe nerve pain.

**percutaneous:** through the skin.

**radiofrequency:** radiant energy of a certain frequency.

**rhizotomy:** section of the nerve roots for the relief of pain.

**trigeminal neuralgia:** a painful disorder of the fifth cranial nerve, called the trigeminal nerve. Irritation of this nerve can cause intense pain that usually affects one side of the face usually in the forehead, cheek, jaw, or teeth.

**trigeminal nerve:** a nerve originating within the brain that supplies feeling and movement to the face. The trigeminal nerve has three divisions: the ophthalmic (V1) division gives sensation to your forehead and eyes; the maxillary (V2) division gives sensation to your cheek, under your eye, around your nose; and the mandibular (V3) division gives sensation to your jaw.

**radiology department:** rooms designed for examining the body by using x-rays.

**stereotaxis:** a precise method for locating deep brain structures by using 3-dimensional coordinates.



updated > 1.2021

reviewed by > Steven Bailey, MD, Mayfield Clinic, Cincinnati, Ohio

Mayfield Certified Health Info materials are written and developed by the Mayfield Clinic. We comply with the HONcode standard for trustworthy health information. This information is not intended to replace the medical advice of your health care provider. © Mayfield Clinic 1998-2021.